

## Physics 115/242

### Numerov method for integrating the one-dimensional Schrödinger equation.

Peter Young

The one-dimensional time-independent Schrödinger equation is

$$-\frac{\hbar^2}{2m} \frac{d^2\psi}{dx^2} + V(x)\psi(x) = E\psi(x), \quad (1)$$

where  $\psi(x)$  is the wavefunction,  $V(x)$  is the potential energy,  $m$  is the mass, and  $\hbar$  is Planck's constant divided by  $2\pi$ . This is an eigenvalue problem since one can only find a solution which vanishes at  $\pm\infty$  (the boundary conditions) for certain discrete values of  $E$ .

In order to find the energy eigenvalues, we need to be able to integrate the equation with respect to  $x$ , for a given value of  $E$ , starting at  $x = x_0$ , say, with some specified values for  $x = x_0$  and  $x = x_1 = x_0 + h$ , where  $h$  is the step interval. Using the notation  $x_n = x_0 + nh$  and  $\psi_n \equiv \psi(x_n)$ , we have to solve for  $\psi_2, \psi_3, \dots$ , given  $\psi_0$  and  $\psi_1$ . Having solved the equation for a given value of  $E$  we need to vary  $E$  until we find a solution which satisfies the boundary conditions, which requires re-solving the equation for each value of  $E$ . We will discuss this aspect of the problem, using what is called the “shooting method”, in more detail in class.

Here we focus on the problem of integrating the equation for a *given* value of  $E$ . One method would be to use 4-th order Runge-Kutta (RK4), since it is quite accurate. RK4 involves writing Schrödinger's equation, which is second order, as two first order equations:

$$\begin{aligned} \frac{d\psi}{dx} &= \phi(x) \\ \frac{d\phi}{dx} &= -k^2(x)\psi(x), \end{aligned} \quad (2)$$

where

$$k^2(x) = \frac{2m}{\hbar^2}(E - V(x)). \quad (3)$$

You will recall that this a fourth order method, i.e. the error is proportional to  $h^4$ .

An alternative, is to leave the Schrödinger equation as one second order equation,

$$\boxed{\left(\frac{d^2\psi}{dx^2} + k^2(x)\right)\psi(x) = 0,} \quad (4)$$

and take advantage of its particular structure (it is linear in  $\psi$  and there is no term involving the first derivative.) A suitable algorithm for this type of problem is the *Numerov* algorithm, which is simpler than RK4 and is one one higher order (fifth).

We now describe the Numerov method (see also Landau and Páez). A Taylor series for  $\psi(x+h)$  gives

$$\psi(x+h) = \psi(x) + h\psi'(x) + \frac{h^2}{2}\psi^{(2)}(x) + \frac{h^3}{6}\psi^{(3)}(x) + \frac{h^4}{24}\psi^{(4)}(x) + \dots \quad (5)$$

Adding this to the series for  $\psi(x-h)$  all the odd powers of  $h$  vanish:

$$\psi(x+h) + \psi(x-h) = 2\psi(x) + h^2\psi^{(2)}(x) + \frac{h^4}{12}\psi^{(4)}(x) + O(h^6). \quad (6)$$

We can therefore write the second derivative which occurs in the Schrödinger equation, Eq. (4), as

$$\psi^{(2)}(x) = \frac{\psi(x+h) + \psi(x-h) - 2\psi(x)}{h^2} - \frac{h^2}{12}\psi^{(4)}(x) + O(h^4). \quad (7)$$

We would like to evaluate the term involving the 4th derivative. To do so, we act on Eq. (4) with  $1 + (h^2/12)d^2/dx^2$ , which gives

$$\psi^{(2)}(x) + \frac{h^2}{12}\psi^{(4)}(x) + k^2(x)\psi(x) + \frac{h^2}{12}\frac{d^2}{dx^2} [k^2(x)\psi(x)] = 0. \quad (8)$$

Substituting for  $\psi^{(2)}(x) + \frac{h^2}{12}\psi^{(4)}(x)$  from Eq. (8) into Eq. (7) gives

$$\psi(x+h) + \psi(x-h) - 2\psi(x) + h^2k^2(x)\psi(x) + \frac{h^4}{12}\frac{d^2}{dx^2} [k^2(x)\psi(x)] + O(h^6) = 0. \quad (9)$$

We evaluate  $\frac{d^2}{dx^2} [k^2(x)\psi(x)]$  by using an elementary difference formula (this has an error  $O(h^2)$  but is accurate enough because this term is already multiplied by  $h^4$  in Eq. (9)):

$$\frac{d^2}{dx^2} [k^2(x)\psi(x)] \simeq \frac{k^2(x+h)\psi(x+h) + k^2(x-h)\psi(x-h) - 2k^2(x)\psi(x)}{h^2}. \quad (10)$$

Substituting Eq. (10) into Eq. (9) and rearranging we get the Numerov algorithm for one time step:

$$\psi(x+h) = \frac{2\left(1 - \frac{5}{12}h^2k^2(x)\right)\psi(x) - \left(1 + \frac{1}{12}h^2k^2(x-h)\right)\psi(x-h)}{1 + \frac{1}{12}h^2k^2(x+h)} + O(h^6). \quad (11)$$

Setting  $x = x_n \equiv x_0 + nh$ , and defining  $k_n \equiv k(x_n)$ , this can be written more tidily as

$$\boxed{\psi_{n+1} = \frac{2\left(1 - \frac{5}{12}h^2k_n^2\right)\psi_n - \left(1 + \frac{1}{12}h^2k_{n-1}^2\right)\psi_{n-1}}{1 + \frac{1}{12}h^2k_{n+1}^2}}, \quad (12)$$

with an error of order  $h^6$ . The Numerov method, Eq. (12), can be used to determine  $\psi_n$  for  $n = 2, 3, 4, \dots$ , given two initial values,  $\psi_0$  and  $\psi_1$ .

The error in one time step is  $O(h^6)$ . However, as we have also discussed in other contexts, the number of steps needed to integrate over a *fixed* range of  $x$  from  $x_0$  to  $x_0 + \Delta x$ , say, is  $\Delta x/h$ . The errors in each step can add up and so the total error in the Numerov method is  $O(h^5)$ , i.e. it is a 5-th order method, one higher than RK4. However, there can be problems with roundoff errors in using Eq. (12) so make sure you use double precision arithmetic.