








Quantum many-body physics near decay thresholds

Alexander Volya
Florida State University

Supported by the US Department of
Energy Award number: DE-SC0009883




Threshold discontinuity





$$H'(\epsilon) = \int_0^\infty d\epsilon' A^*(\epsilon') \frac{1}{\epsilon - \epsilon' + i0} A(\epsilon')$$

Effect of weak binding



Effect of weak binding





C. R. Hoffman, B. P. Kay, and J. P. Schiffer Phys. Rev. C 89, 061305(R)

B. P. Kay, C. R. Hoffman, and A. O. Macchiavelli Phys. Rev. Lett. 119, 182502

Wave function realignment Superradiance

- Model $\mathcal{H}$



$$\mathcal{H}(E) = \begin{pmatrix} \epsilon_1 - \frac{i}{2}\gamma_1 & v - \frac{i}{2}A_1A_2 \\ v - \frac{i}{2}A_1A_2 & \epsilon_2 - \frac{i}{2}\gamma_2 \end{pmatrix}$$



Example of interacting resonances

^{13}C , ^{13}N , ^{11}Li

$$\mathcal{H} = H^0 + V - iW/2$$




Imaginary W

$W \neq 0$

$W=0$



^{13}C interference of levels



K. Hanselman *Direct Reactions And Decay Correlations Studying The Proton-Unbound Superradiant State In ^{13}N* , Thesis, Florida State University 2022

^{13}C interference of levels

	Exp. <u>Energy</u>	Exp. <u>Width</u>	SM <u>Width</u>	CSM <u>Width</u>
$3/2^+(1)$	7.686(6)	0.070(5)	0.809	0.094
$3/2^+(2)$	8.2(1)	1.1(3)	0.369	0.984

K. Kravvaris and A. Volya, "Quest for superradiance in atomic nuclei," AIP Conf. Proc., vol. 1912, no. 1, p. 020010, 2017, doi: 10.1063/1.5016135.

K. Hanselman *Direct Reactions And Decay Correlations Studying The Proton-Unbound Superradiant State In ^{13}N* , Thesis, Florida State University 2022

Wave function realignment

Superradiance

$$H = \begin{pmatrix} \epsilon - \frac{i}{2}\Gamma & v \\ v & 0 \end{pmatrix} = H_0 - \frac{i\Gamma}{2} A^\dagger A \quad A = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

Stationary system $\Gamma = 0$

Energies
$$E_{1,2} = \frac{1}{2} \left(\epsilon \pm \sqrt{\epsilon^2 + 4v^2} \right)$$

Spectroscopic Factors
$$\text{SF}_{1,2} = \frac{1}{2} \left(1 \pm \frac{\epsilon}{\sqrt{\epsilon^2 + 4v^2}} \right)$$

Observing superradiance

$$H = \begin{pmatrix} \epsilon - \frac{i}{2}\Gamma & v \\ v & 0 \end{pmatrix} = H_0 - \frac{i\Gamma}{2} A^\dagger A \quad A = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

Energies

$$\mathcal{E}_{1,2} = \frac{1}{2} \left(\epsilon - \frac{i}{2}\Gamma \pm \sqrt{\left(\epsilon - \frac{i}{2}\Gamma \right)^2 + 4v^2} \right)$$


Width

$$\Gamma_{1,2} = -2 \operatorname{Im}(\mathcal{E}_{1,2})$$


Spectroscopic Factors

$$\text{SF}_{1,2} = \Gamma_{1,2}/\Gamma.$$


Observing superradiance




Spectroscopic factor for superradiant state




Clustering in A=18 mirror systems



Clustering in A=18 mirror systems







Isospin symmetry



	^{18}Ne			^{18}O		
J	E (MeV)	Γ (keV)	SF	E (MeV)	Γ (keV)	SF
1-	9.08(1)	357	0.21(1)	9.19(2)	200	0.20(1)
1-	9.57(1)	1062	0.51(5)	9.76(2)	630	0.46(4)
1-	10.58(4)	416	0.15(5)	10.8(3)	630	0.29(4)
1-	13.730(2)	780	0.2(1)	14.3(3)	400	0.10(4)
2+	9.19(3)	265	0.21(2)	9.79(6)	90	0.10(3)
2+	10.94(6)	1302	0.52(3)	12.21(8)	1000	0.37(9)
2+	13.4 (2)	1755	0.45(8)	12.8(3)	4800	1.56(13)
2+	16.9(2)	1515	0.3(2)			
3-				8.29(6)	2.9	0.18(1)
3-	8.77(8)	419	1.0(4)	9.35(2)	110	0.48(13)
3-	11.0(1)	497	0.28(7)	11.95(1)	300	0.17(2)
3-	12.7(2)	2025	0.7(2)	12.98(4)	770	0.32(5)
3-	14.8(2)	3967	1.0(2)	14.0(2)	2100	0.7(1)
4+	8.16	31	0.8(3)	7.11*		
4+	13.3(3)	845	0.37(4)	13.46(2)	210	0.12(1)
4+	14.15(21)	375	0.14(10)	14.77(5)	680	0.28(2)
5-	11.31(4)	15	0.03(2)	11.63(1)	30	0.13(1)
5-	12.9(2)	532	0.48(12)	13.08(1)	120	0.17(1)
5-	13.79(8)	219	0.14(10)	14.1(1)	260	0.23(2)
5-	14.6(7)	521	0.27(20)	14.7(1)	230	0.16(6)
6+	11.8(2)	54	0.30(7)	11.69(5)	12	0.23(1)
6+	12.4(2)	167	0.56(26)	12.57(1)	50	0.38(8)

M. Barbui *et al.*, “ α -cluster structure of ^{18}Ne ,”
Phys. Rev. C, vol. 106, no. 5, p. 054310, Nov.
2022, doi: [10.1103/PhysRevC.106.054310](https://doi.org/10.1103/PhysRevC.106.054310).

	^{18}Ne			^{18}O		
J	E (MeV)	Γ (keV)	SF	E (MeV)	Γ (keV)	SF
1-	9.08(1)	357	0.21(1)	9.19(2)	200	0.20(1)
1-	9.57(1)	1062	0.51(5)	9.76(2)	620	0.46(1)




5-	14.6(7)	521	0.27(20)	14.7(1)	230	0.16(6)
6+	11.8(2)	54	0.30(7)	11.69(5)	12	0.23(1)
6+	12.4(2)	167	0.56(26)	12.57(1)	50	0.38(8)


M. Barbui *et al.*, “ α -cluster structure of ^{18}Ne ,” *Phys. Rev. C*, vol. 106, no. 5, p. 054310, Nov. 2022, doi: [10.1103/PhysRevC.106.054310](https://doi.org/10.1103/PhysRevC.106.054310).

	^{18}Ne			^{18}O		
J	E (MeV)	Γ (keV)	SF	E (MeV)	Γ (keV)	SF
1-	9.08(1)	357	0.21(1)	9.19(2)	200	0.20(1)
1-	9.57(1)	1062	0.51(5)	9.76(2)	630	0.46(4)
1-	10.58(4)	416	0.15(5)	10.8(3)	630	0.29(4)
1-	13.730(2)	780	0.2(1)	14.3(3)	400	0.10(4)
2+	9.19(3)	265	0.21(2)	9.79(6)	90	0.10(3)
2+	10.94(6)	1302	0.52(3)	12.21(8)	1000	0.37(9)
2+	13.4 (2)	1755	0.45(8)	12.8(3)	4800	1.56(13)
2+	16.9(2)	1515	0.3(2)			
3-				8.29(6)	2.9	0.18(1)
3-	8.77(8)	419	1.0(4)	9.35(2)	110	0.48(13)
3-	11.0(1)	497	0.28(7)	11.95(1)	300	0.17(2)
3-	12.7(2)	2025	0.7(2)	12.98(4)	770	0.32(5)
3-	14.8(2)	3967	1.0(2)	14.0(2)	2100	0.7(1)
4+	8.16	31	0.8(3)	7.11*		
4+	13.3(3)	845	0.37(4)	13.46(2)	210	0.12(1)
4+	14.15(21)	375	0.14(10)	14.77(5)	680	0.28(2)
5-	11.31(4)	15	0.03(2)	11.63(1)	30	0.13(1)
5-	12.9(2)	532	0.48(12)	13.08(1)	120	0.17(1)
5-	13.79(8)	219	0.14(10)	14.1(1)	260	0.23(2)
5-	14.6(7)	521	0.27(20)	14.7(1)	230	0.16(6)
6+	11.8(2)	54	0.30(7)	11.69(5)	12	0.23(1)
6+	12.4(2)	167	0.56(26)	12.57(1)	50	0.38(8)


M. Barbui *et al.*, “ α -cluster structure of ^{18}Ne ,”
Phys. Rev. C, vol. 106, no. 5, p. 054310, Nov.
2022, doi: [10.1103/PhysRevC.106.054310](https://doi.org/10.1103/PhysRevC.106.054310).



Clustering and continuum



Clustering in light nuclei



- Clusters mix structure and reactions
- Strength distribution


The Nuclear Shell Model

$$H = \frac{\mathbf{p}^2}{2MA} + H_{\text{int}} \quad H(\omega_0) = H + \frac{AM\omega_0^2}{2} \mathbf{R}^2 = H_{\text{cm}}(\omega_0) + H_{\text{int}}$$

$$H_{\text{cm}}(\omega_0) = \frac{\mathbf{p}^2}{2MA} + \frac{AM\omega_0^2}{2} \mathbf{R}^2 \quad \Psi_{n\ell m} = \phi_{n\ell m}(\mathbf{R}) \Psi'$$


$$E(\omega_0) = \hbar\omega_0(N_{\text{cm}} + 3/2) + E'$$

$$N_{\text{cm}} = 2n + \ell$$



Translational invariance and Center of Mass (CM)

Shell model, Glockner-Lawson procedure



Controlling CM with operator

$\mathbf{R}$

Control only
CM quanta

$$D_\mu = \sqrt{\frac{4\pi}{3}} R_\mu \quad R_\mu = \sqrt{\frac{\hbar}{2Am\omega}} (\mathcal{B}_\mu^\dagger + \mathcal{B}_\mu)$$


K. Kravvaris and A. Volya, “Study of nuclear clustering from an ab initio perspective,” *Phys. Rev. Lett.*, vol. 119, no. 6, p. 062501, 2017.

K. Kravvaris and A. Volya, “Clustering in structure and reactions using configuration interaction techniques,” *Phys. Rev. C*, vol. 100, no. 3, p. 034321, Sep. 2019, doi: [10.1103/PhysRevC.100.034321](https://doi.org/10.1103/PhysRevC.100.034321).

Configuration interaction approach and clustering

Traditional shell model configuration m-scheme

$$|\Psi\rangle = \Psi^\dagger |0\rangle \sim a_1^\dagger a_2^\dagger \dots a_A^\dagger |0\rangle$$




$|\Psi\rangle$

+

Cluster configuration


$$|\text{channel}\rangle \sim |\Phi\Psi_D\rangle \equiv \Phi^\dagger \Psi_D^\dagger |0\rangle$$



$\Phi^\dagger |\Psi_D\rangle$

Clustering reaction basis channel

(basis states for clustering)



$$\Psi = \phi_{000}(\mathbf{R}) \Psi'$$

Boost

$$\Psi_{n\ell m} = \phi_{n\ell m}(\mathbf{R}) \Psi'$$

CM-Recouple


$$\Phi_{n\ell m} = \mathcal{A} \left\{ \phi_{000}(\mathbf{R}) \phi_{n\ell m}(\boldsymbol{\rho}) \Psi'^{(1)} \Psi'^{(2)} \right\}$$

$$\Phi_{n\ell}^\dagger = \sum_{\substack{n_1 \ell_1 \\ n_2 \ell_2}} \mathcal{M}_{n_1 \ell_1 n_2 \ell_2}^{n\ell 00; \ell} \left[\Psi_{n_1 \ell_1 m_1}^\dagger \times \Psi_{n_2 \ell_2 m_2}^\dagger \right]_\ell$$


Resonating group method and reactions

$$\sum_n \mathcal{H}_{nn'}^{(\ell)} \chi_{n'} = E \sum_n \mathcal{N}_{nn'}^{(\ell)} \chi_{n'}$$

$$\begin{pmatrix} \mathcal{H}_{00} & \dots & \mathcal{H}_{0n} & 0 & 0 & 0 \\ \vdots & \ddots & \vdots & 0 & \vdots & \vdots \\ \mathcal{H}_{n0} & \dots & \mathcal{H}_{nn} & T_{nn+1} & 0 & \vdots \\ 0 & 0 & T_{n+1n} & T_{n+1n+1} & T_{n+1n+2} & 0 \\ 0 & \dots & 0 & T_{n+2n+1} & T_{n+2n+2} & \ddots \\ 0 & \dots & \dots & 0 & \ddots & \ddots \end{pmatrix} \begin{pmatrix} \chi_0 \\ \vdots \\ \chi_n \\ x_{n+1} \\ x_{n+2} \\ \vdots \end{pmatrix} = E \begin{pmatrix} \mathcal{N}_{00} & \dots & \mathcal{N}_{0n} & 0 & 0 & 0 \\ \vdots & \ddots & \vdots & 0 & \vdots & \vdots \\ \mathcal{N}_{n0} & \dots & \mathcal{N}_{nn} & 0 & 0 & \vdots \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & \dots & 0 & 0 & 1 & 0 \\ 0 & \dots & \dots & 0 & 0 & \ddots \end{pmatrix} \begin{pmatrix} \chi_0 \\ \vdots \\ \chi_n \\ x_{n+1} \\ x_{n+2} \\ \vdots \end{pmatrix}$$




Asymptotic solution with phase shift




J-matrix (or HORSE) method: J. M. Bang, Annals of Physics **280**, 299 (2000)
Experimental data: Phys. Rev. 168, 1114 (1968); Nucl. Phys. **A287**, 317 (1977)

alpha+alpha scattering phase shifts




Experimental data from S. A. Afzal, A. A. Z. Ahmad, and S. Ali, Rev. Mod. Phys. 41, 247 (1969).

Asymptotic Normalization and blocking




ANC and role of blocking




Spectroscopic amplitudes

parent	J^π	channel	$ \langle \Psi \mathcal{F}_e \rangle $
$^8\text{Be}[4]$	0^+	$\alpha[0] + \alpha[0]$	0.905
$^8\text{Be}[4]$	2^+	$\alpha[0] + \alpha[0]$	0.898
$^8\text{Be}[4]$	4^+	$\alpha[0] + \alpha[0]$	0.874
$^8\text{Be}[4]$	0^+	$\alpha[2] + \alpha[2]$	0.961
$^8\text{Be}[4]$	2^+	$\alpha[2] + \alpha[2]$	0.957
$^8\text{Be}[4]$	4^+	$\alpha[2] + \alpha[2]$	0.943
$^{10}\text{Be}[4]$	0^+	$^6\text{He}[0] + \alpha[0]$	0.844
$^{10}\text{Be}[4]$	0^+	$^6\text{He}[4] + \alpha[0]$	0.820
$^{10}\text{Be}[4]$	2^+	$^6\text{He}[0] + \alpha[0]$	0.834
$^{10}\text{Be}[4]$	2^+	$^6\text{He}[4] + \alpha[0]$	0.796
$^{12}\text{C}[4]$	0_1^+	$\alpha[0] + \alpha[0] + \alpha[0]$	0.841
$^{12}\text{C}[4]$	0_2^+	$\alpha[0] + \alpha[0] + \alpha[0]$	0.229




Spectroscopic amplitudes.


Ttriple-alpha RGM




$N_{\max}(\text{rel})=12$



parent	channel	overlap
$^{12}\text{C}[4](0_1^+)$	$\alpha[0] + \alpha[0] + \alpha[0]$	0.841
$^{12}\text{C}[4](0_2^+)$	$\alpha[0] + \alpha[0] + \alpha[0]$	0.229




Searching for clustering strength



α -spectroscopic factors for states in ^{34}Ar calculated using the shell model

A. M. Long *et al.*, “alpha-unbound levels in Ar-^{34} from $\text{Ar-}^{36}(\text{p},\text{t})\text{Ar-}^{34}$ reaction measurements and implications for the astrophysical $\text{S-}^{30}(\alpha,\text{p})\text{Cl-}^{33}$ reaction rate,” *Phys. Rev. C*, vol. 97, no. 5, p. 054613, May 2018, doi: [10.1103/PhysRevC.97.054613](https://doi.org/10.1103/PhysRevC.97.054613).


Searching for clustering strength



Distribution of dynamic spectroscopic factors for $^{20}\text{Ne} \rightarrow ^{16}\text{O}(\text{g.s.}) + \alpha$. The dashed lines correspond to the RGM energies for each decay channel.


Channel coupling in ^{18}O

$I=1$ channel




Channel coupling in ^{18}O

$l=1$ channel





Channel coupling in ^{18}O



$l=1$ channel



Channel coupling in ^{18}O



^{11}Be beta-delayed proton decay




Observed half-life (???)

$$t_{\text{Be} \rightarrow \beta p} \approx 1 \times 10^6 \text{ s.}$$

$$\underline{1/2^+ \quad 9.820}$$

$$\alpha + \text{}^7_3\text{Li}$$

^{11}Be beta-delayed proton decay



E. Lopez-Saavedra *et al.*, *Phys. Rev. Lett.*, vol. 129, no. 1, p. 012502, Jun. 2022, doi: [10.1103/PhysRevLett.129.012502](https://doi.org/10.1103/PhysRevLett.129.012502).

Y. Ayyad *et al.*, *Phys. Rev. Lett.*, vol. 129, no. 1, p. 012501, Jun. 2022, doi: [10.1103/PhysRevLett.129.012501](https://doi.org/10.1103/PhysRevLett.129.012501).

Observed half-life (???)

$$t_{\text{Be} \rightarrow \beta p} \approx 1 \times 10^6 \text{ s.}$$

$\alpha + ^7_3\text{Li}$

Notes on beta-decay rates

$$ft = \frac{\mathcal{T}}{B(F) + \lambda_A^2 B(GT)} \quad \mathcal{T} = 6145 \text{ s},$$

$$\lambda_A = 1.27$$


Neutron decay $n \rightarrow p + e^- + \bar{\nu}_e$ $B(F) = 1$ $B(GT) = 3$


$$f(Q) = 1.715$$

Sequential beta decay


For sequential decay


$$\mathcal{F}(Q) = \int_0^Q \frac{d\epsilon}{2\pi} \frac{f(\epsilon) \gamma_2(Q - \epsilon)}{(\epsilon + E_2)^2 + \Gamma_2^2(Q - \epsilon)/2}$$






$$t_{\text{Be} \rightarrow \beta p} \approx 1 \times 10^6 \text{ s.}$$





1/2+ states in 11B

	Theory (FSU)			Experiment	
J	E(MeV)	log(ft)	SF(p)	E(MeV)	SF(p)
1/2+(1)	5.709	5.5	0.262	6.792	
1/2+(2)	10.545	3.4	0.117	9.820	
1/2+(3)	11.952	3.5	0.134	11.44	0.27(6)
1/2+(4) T=3/2	12.181		0.274	12.554	
1/2+(5)	12.827	4.0	0.028		
1/2+(6)	14.105	5.4	0.001		






Observations

- The proton resonance is likely $1/2^+(3)$, and it is lowered (predicted 12.2 MeV, observed at 11.44 MeV)
- The proton SF is large, while there is no alpha decay?
- The strong GT transition is difficult to explain.



Time dependence of decay

Winter, *Phys. Rev.*, **123**,1503 1961.



Internal dynamics in decaying system

Winter's model



M. Peshkin, A. Volya, and V. Zelevinsky, “Non-exponential and oscillatory decays in quantum mechanics,” *EPL*, vol. 107, no. 4, p. 40001, 2014.

Time-dependent Continuum Shell Model Approach

- Expand Using evolution operator in Chebyshev polynomials

$$\exp(-iHt) = \sum_{n=0}^{\infty} (-i)^n (2 - \delta_{n0}) J_n(t) T_n(H)$$

- Chebyshev polynomial $T_n[\cos(\theta)] = \cos(n\theta)$
- Use iterative relation and matrix-vector multiplication to generate

$$|\lambda_n\rangle = T_n(H)|\lambda\rangle$$

$$|\lambda_0\rangle = |\lambda\rangle, \quad |\lambda_1\rangle = H|\lambda\rangle \quad |\lambda_{n+1}\rangle = 2H|\lambda_n\rangle - |\lambda_{n-1}\rangle$$


$$\langle\lambda'|T_{n+m}(H)|\lambda\rangle = 2\langle\lambda'_m|\lambda_n\rangle - \langle\lambda'|\lambda_{n-m}\rangle, \quad n \geq m$$

- Use FFT to find return to energy representation

Probing the Non-exponential Decay Regime in Open Quantum Systems

- Broad threshold resonance (^9N , ^9He)
 - Pronounced non-exponentiality
 - Very short half-life
- Three-body decay (^6Be , ^{13}Li , ^{16}Be)
 - Nucleon-nucleon correlations
 - Energy dependence
- Overlapping resonances (^{13}C , ^{13}N)
 - Interference, pronounced non-exponentiality
 - Superradiance


Time-dependent picture



$$\mathcal{G} = \frac{1}{E - E_o + i/2 \Gamma(E)}$$

$$\Gamma(E) \propto \sqrt{E}$$

Power-law remote decay rate!



Probing the Non-exponential Decay Regime in Open Quantum Systems

- Broad threshold resonance (^9N , ^9He)
 - Pronounced non-exponentiality
 - Very short half-life
- Three-body decay (^6Be , ^{13}Li , ^{16}Be)
 - Nucleon-nucleon correlations
 - Energy dependence
- Overlapping resonances (^{13}C , ^{13}N)
 - Interference, pronounced non-exponentiality
 - Superradiance

Probing the Non-exponential Decay Regime in Open Quantum Systems



- Broad threshold resonance (^9N , ^9He)
 - Pronounced non-exponentiality
 - Very short half-life
- Three-body decay (^6Be , ^{13}Li , ^{16}Be)
 - Nucleon-nucleon correlations
 - Energy dependence
- Overlapping resonances (^{13}C , ^{13}N)
 - Interference, pronounced non-exponentiality
 - Superradiance

Two-level system

$$\mathcal{H} = \begin{pmatrix} \epsilon_1 - (i/2)\Gamma_1 & v - (i/2)A_1A_2 \\ v - (i/2)A_1A_2 & \epsilon_2 - (i/2)\Gamma_2 \end{pmatrix}$$

$$\Gamma_1 = A_1^2, \quad \Gamma_2 = A_2^2,$$

$$S(t) = |\langle \Psi(0) | \Psi(t) \rangle|^2$$



$$\mathcal{H} = H^0 + V - iW/2$$

Real V

$V=0$




$V \neq 0$

Imaginary W


$W \neq 0$



$W=0$




Wigner cusps, unitarity ...




$$\Phi(X, x) = \frac{e^{iK_n X}}{\sqrt{|K_n|}} \psi_n(x) + \sum_{m=0}^{\infty} \frac{R_{mn}}{\sqrt{|K_m|}} e^{-iK_m X} \psi_m(x),$$

$$\Phi(X, x) = 0 \text{ at } x_2 = 0.$$





Two-nucleon system with continuum finite square well binding


In the exact solution virtual amplitudes are exponentially large.





N. Ahsan and A. Volya, "Quantum tunneling and scattering of a composite object reexamined," *Phys. Rev. C*, vol. 82, no. 6, p. 064607, 2010.

Results: scattering off an infinite wall







Resonant tunneling of composite objects



Enhanced tunneling probability for composite objects



Enhancement of tunneling probability.



^6He

References:

- [1] M. Peshkin, A. Volya, and V. Zelevinsky, “Non-exponential and oscillatory decays in quantum mechanics,” EPL, vol. 107, no. 4, p. 40001, 2014.
- [2] V. Zelevinsky and A. Volya, Mesoscopic Nuclear Physics: From Nucleus to Quantum Chaos to Quantum Signal Transmission. WORLD SCIENTIFIC, 2023. doi: 10.1142/13049.
- [3] K. K. W. Ma, A. Volya, and K. Yang, “Eigenstate thermalization and disappearance of quantum many-body scar states in weakly interacting fermion systems,” Phys. Rev. B, vol. 106, no. 21, p. 214313, Dec. 2022, doi: 10.1103/PhysRevB.106.214313.
- [4] A. Volya, M. Barbui, V. Z. Goldberg, and G. V. Rogachev, “Superradiance in alpha clustered mirror nuclei,” Commun Phys, vol. 5, no. 1, Art. no. 1, Dec. 2022, doi: 10.1038/s42005-022-01105-9.
- [5] E. Lopez-Saavedra et al., “Observation of a near-threshold proton resonance in ^{11}B ,” Phys. Rev. Lett., vol. 129, no. 1, p. 012502, Jun. 2022, doi: 10.1103/PhysRevLett.129.012502.
- [6] M. Barbui et al., “ α -cluster structure of ^{18}Ne ,” Phys. Rev. C, vol. 106, no. 5, p. 054310, Nov. 2022, doi: 10.1103/PhysRevC.106.054310.
- [7] A. Volya and V. Zelevinsky, “Time-dependent relaxation of observables in complex quantum systems,” J. Phys. Complex., vol. 1, no. 2, p. 025007, 2020, doi: 10.1088/2632-072x/ab79bc.
- [8] A. Volya, H. A. Weidenmüller, and V. Zelevinsky, “Neutron resonance widths and the porter-thomas distribution,” Phys.Rev.Lett., vol. 115, no. 5, p. 052501, Jul. 2015, doi: 10.1103/PhysRevLett.115.052501.
- [9] A. Volya, “Computational approaches to many-body dynamics of unstable nuclear systems,” in Nuclear theory in the super- computing era — 2014, A. M. Shirokov and A. I. Mazur, Eds., 2015, Ju 2014, p. 21.
- [10] A. Volya, “Physics of unstable nuclei: from structure to sequential decays,” EPJ Web of Conf., vol. 38, p. 03003, 2012.
- [11] A. Volya, “Porter-Thomas distribution in unstable many-body systems,” Phys. Rev. C, vol. 83, no. 4, p. 044312, 2011, doi: 10.1103/PhysRevC.83.044312.
- [12] A. Volya, “Time-dependent approach to the continuum shell model,” Phys. Rev. C, vol. 79, no. 4, p. 044308, 2009, doi: 10.1103/PhysRevC.79.044308.
- [13] A. Volya and V. Zelevinsky, “Continuum shell model,” Phys. Rev. C, vol. 74, no. 6, p. 064314, 2006.
- [14] A. Volya and V. Zelevinsky, “Non-Hermitian effective Hamiltonian and continuum shell model,” Phys. Rev. C, vol. 67, no. 5, p. 054322, May 2003.
- [15] S. M. Wang, W. Nazarewicz, A. Volya, and Y. G. Ma, “Probing the Non-exponential Decay Regime in Open Quantum Systems.” arXiv, Nov. 21, 2022. doi: [10.48550/arXiv.2211.11619](https://arxiv.org/abs/10.48550/arXiv.2211.11619).
- [16] N. Ahsan and A. Volya, “Quantum tunneling and scattering of a composite object reexamined,” Phys. Rev. C, vol. 82, no. 6, p. 064607, 2010.
- [17] A. M. Long *et al.*, “alpha-unbound levels in Ar-34 from Ar-36(p,t)Ar-34 reaction measurements and implications for the astrophysical S-30(alpha, p)Cl-33 reaction rate,” Phys. Rev. C, vol. 97, no. 5, p. 054613, May 2018, doi: [10.1103/PhysRevC.97.054613](https://doi.org/10.1103/PhysRevC.97.054613).

Funding Acknowledgements:

Funding: U.S. DOE contract DE-SC0009883.